

# Effects of Cigarette Smoking on Serum Lipids among Priests in Bangkok

Pairat Saengdith, MD\*

\*Department of Medicine, Priest Hospital

**Objectives:** To determine the effects of cigarette smoking on the level of serum cholesterol and triglyceride among priests in Bangkok area.

**Materials and Method:** Serum total cholesterol and triglyceride were evaluated in the process of health check-up among 401 priests in Bangkok and a questionnaire about cigarette smoking was given.

**Results:** From the 401 priests, 76.3% of priests were more than 40 years old. The mean age of priests was  $53.68 \pm 15.01$  years. The mean cholesterol among non-smokers, ex-smokers, current smokers and all respondents were  $230.08 \pm 48.22$ ,  $223.58 \pm 42.80$ ,  $230.08 \pm 48.22$ , and  $226.65 \pm 46.58$  respectively. The mean triglyceride among non-smokers, ex-smokers, current smokers and all respondents were  $150.05 \pm 78.23$ ,  $148.47 \pm 69.80$ ,  $180.01 \pm 111.99$ , and  $154.87 \pm 83.71$ , respectively. In comparison among groups, there was no significantly statistical difference in cholesterol level ( $p = 0.22$ ), but there was significantly statistical difference in triglyceride level ( $p = 0.02$ ). The means of triglyceride between non-smokers and current smokers, and between ex-smokers and current smokers were significantly statistical different ( $p = 0.026$ ,  $0.043$ , respectively).

**Conclusion:** The results suggest that cigarette smoking increases serum triglyceride level in the priests but does not affect serum total cholesterol level. The priests who usually smoke should be checked for triglyceride for early intervention of coronary heart disease and other hypertriglyceridemia-induced diseases.

**Keywords:** Smoking, Lipid, Cholesterol, Triglyceride, Priests, Bangkok

*J Med Assoc Thai 2008; 91 (Suppl 1): S41-4*

*Full text. e-Journal:* <http://www.medassocthai.org/journal>

Cigarette smoking is one of the major risk factors for cardiovascular disease. Smokers have a higher risk of coronary artery disease than non-smokers. Several possible explanations have been offered for this association, including altered blood coagulation, impaired integrity of the arterial wall, and changes in blood lipids<sup>(1)</sup>. Several studies have shown an association between cigarette smoking and altered serum lipids. In conclusion, the studies revealed that cigarette smokers have lower high-density lipoprotein (HDL) cholesterol levels and higher levels of total cholesterol, low-density lipoprotein (LDL) cholesterol, and triglyceride than non-smokers<sup>(2,3)</sup>. It is evident that smoking causes endothelial dysfunction, decreased HDL cholesterol levels, hypertriglyceridemia and

increased oxidation of LDL cholesterol and platelet activation leading to a prothrombotic state<sup>(4)</sup>.

In Thailand, there were some studies involving the association between smoking and blood lipids. The study of 501 clerks of Government Savings Bank showed that smoking and alcohol consumption were associated with hypertriglyceridemia and decreased HDL cholesterol, but were not associated with total cholesterol level<sup>(5)</sup>. The study of 80 male Bangkok Metropolitan Waterworks employees showed that smokers had significant lower levels of HDL cholesterol than non-smokers<sup>(6)</sup>. Another study of 133 clients for health check-up at Rajavithi Hospital showed that smoking affected decreasing of HDL cholesterol<sup>(7)</sup>. Priests is a special group of population, i.e., all of them are male, the age is 20 years and above, they eat only 2 meals per day, their daily life usually does not involve with heavy exercise, they do not drink alcohol and the prevalence of smoking among them is around 25 %<sup>(8)</sup>.

Correspondence to: Saengdith P. Priest Hospital, 445 Sri Ayudhya street, Bangkok 10400 Thailand. Phone: 0-2640-9537, Fax: 0-2354-4287.

Priest Hospital started health check-up of priests in Bangkok in 2006. It is interesting to study the effects of smoking on their serum lipid for possible association among these variables and to get some clues to decrease coronary heart disease (CHD) among them.

### Objective

To determine the effects of cigarette smoking on the level of serum total cholesterol and triglyceride in priests from Bangkok area.

### Material and Method

Four hundred and one priests in Bangkok who participated in health check-up by Priest Hospital in 2006 were subjects of this study. Serum total cholesterol and triglyceride were evaluated from venous blood and a questionnaire about cigarette smoking was given. The subjects were classified according to smoking statuses into three groups, i.e., non-smokers, ex-smokers, and

current smokers. The data were analyzed by using SPSS version 11 software to determine means, standard deviations, associations between smoking status and blood lipids and comparisons of mean difference of blood lipids between pairs of smoking status.

### Results

Almost all of the respondents were more than 30 years of age, 65% of them were between 31-60 and 76.3% of them were more than 40 years old (Table 1). The mean age was  $53.68 \pm 15.01$  years. The mean cholesterol among non-smokers, ex-smokers, and current smokers were  $230.08 \pm 48.22$ ,  $223.58 \pm 42.80$ , and  $230.08 \pm 48.22$ , respectively. The mean triglyceride among non-smokers, ex-smokers, and current smokers were  $150.05 \pm 78.23$ ,  $148.47 \pm 69.80$ , and  $180.01 \pm 111.99$ , respectively (Table 2). In comparison among groups, there was no significantly statistical difference in cholesterol level ( $p = 0.22$ ), but there was significantly statistical difference in triglyceride level ( $p = 0.02$ ) (Table 2). The means of triglyceride between non-smokers and current smokers, and between ex-smokers and current smokers were significantly statistical different ( $p = 0.026, 0.043$ , respectively (Table 3).

### Discussion

Measurement of blood lipids in the clinical laboratory has become increasingly important because of the predictive association with cardiovascular diseases, especially coronary artery disease<sup>(9-12)</sup>. Lipid profiles in health check-up are usually the evaluation of total cholesterol, triglyceride, HDL cholesterol, and LDL cholesterol. In this study, only total cholesterol and triglyceride were measured for economic purpose. It showed that the mean triglyceride of non-smokers was significantly statistical different from current

**Table 1.** Number of priests classified by age groups

Age group (year)	Number	Percent
Less than 30	9	2.2
31-40	86	21.4
41-50	93	23.2
51-60	82	20.4
More than 60	131	32.7
Total	401	100.0

**Table 2.** Comparisons of means of serum lipid classified by smoking statuses and association between smoking and serum lipid

Lipid type/ Smoking status	n	Mean	SD	F	p-value
Chol Non-smoker	226	230.08	48.22	1.517	0.221
Ex-smoker	105	223.58	42.80		
Cur. smoker	70	220.18	46.26		
Total	401	226.65	46.58		
TG Non-smoker	226	150.05	78.23	3.895	0.021*
Ex-smoker	105	148.47	69.80		
Cur.smoker	70	180.01	111.99		
Total	401	154.87	83.71		

\* significant level at  $p < 0.05$

Chol = Cholesterol TG = Triglyceride

Cur.smoker = Current smoker

**Table 3.** Comparisons of means of triglyceride classified by smoking status (n = 401)

Smoking status		Mean difference	p-value
Non-smoker	Ex-smoker	1.59-29.96	1.0000.026*
	Current smoker		
Ex-smoker	Non-smoker	-1.59	1.000
	Current smoker	-31.55	0.043*
Current smoker	Non-smoker	29.96	0.026*
	Ex-smoker	31.55	0.043*

\* significant level at  $p < 0.05$

smokers and ex-smoker or smoking is associated with significantly higher serum concentrations of triglyceride which is the same as the other studies. However, the result of the association with total cholesterol is different from the other studies<sup>(3,5)</sup>. There are many factors that are involved in the different results, such as, diet, physical activity, body weight and the waist/hip ratio, smoking, alcohol consumption and psychosocial conditions<sup>(13)</sup>. The other factor that affects blood lipids are genetic variation<sup>(14)</sup>. The effect of smoking on the serum lipid levels is also dependent on age and gender<sup>(15)</sup>. The triglyceride level is one of several lipid parameters that can aid prediction of CHD risk. An elevated plasma triglyceride level is strongly associated with an increased risk of CHD<sup>(16-18)</sup>. Raised triglyceride levels can be present in individuals at risk for CHD when the total cholesterol is normal. The role of triglyceride in CHD pathogenesis is thought to involve several direct and indirect mechanisms, such as effects on the metabolism of other lipoproteins, transport proteins, enzymes, and on coagulation and endothelial dysfunction<sup>(19)</sup>. However, not all individuals with raised triglyceride levels have increased risk of CHD. Factors, such as, genetic defects, diet, age, lifestyle, and a range of medical conditions, drug therapy and metabolic disorders, can also affect the triglyceride level<sup>(20)</sup>. Hypertriglyceridemia is frequently associated with other lipid abnormalities and the metabolic syndrome, which are linked to coronary artery disease<sup>(21-23)</sup>. Smoking is also a risk factor for pancreatitis<sup>(20)</sup>. Elevated triglyceride level in priests may be partly caused by eating more sugar than general male population. However, the priests who are smokers should be checked for serum lipids, especially triglyceride for early intervention of CHD and other diseases induced by hypertriglyceridemia.

### **Conclusion**

The results suggest that cigarette smoking increases serum triglyceride level in this group of priests, but does not affect serum total cholesterol level. The priests who usually smoke should be checked for serum lipids, especially triglyceride for early intervention of CHD and other hypertriglyceridemia-induced diseases.

### **Acknowledgement**

The author wish to thank Dr. Sawasdi Thakerngdej, the ex-director of Priest Hospital who initiated the health check-up program for the priests and Dr. Waraporn Bhumisawasdi, Director of Priest Hospital who supported this study.

### **References**

- McGill HC Jr. Potential mechanisms for the augmentation of atherosclerosis and atherosclerotic disease by cigarette smoking. *Prev Med* 1979; 8: 390-403.
- Maeda K, Noguchi Y, Fukui T. The effects of cessation from cigarette smoking on the lipid and lipoprotein profiles: a meta-analysis. *Prev Med* 2003; 37: 283-90.
- Craig WY, Palomaki GE, Haddow JE. Cigarette smoking and serum lipid and lipoprotein concentrations: an analysis of published data. *BMJ* 1989; 298: 784-8.
- Tsiara S, Elisaf M, Mikhailidis DP. Influence of smoking on predictors of vascular disease. *Angiology* 2003; 54: 507-30.
- Chularojanamontri V, Kewsiri D, Chuaprasert S, Chanrit S, Viseshakul D. Effects of cigarette smoking and alcohol consumption on blood lipid variables. *J Med Assoc Thai* 1985; 68: 503-7.
- Chatkaew C. Effect of cigarette smoking on lipid profiles in Metropolitant Waterworks employees [Thesis for Master of Science in Public Health (Nutrition)]. Bangkok: Mahidol University; 1990.
- Narawiwat N. Cigarette smoking and serum cholesterol, triglyceride, blood pressure, plasma glucose and kidney function. *Reg 4 Med J* 2003; 22: 177-81.
- Charoenkha N, Kangsakulniti N, Kengkarnpanich, Kusolwisitkul W, Pichainarong N, Silapasawan , et al. Smoking of priests in Thailand. Bangkok: Report of Faculty of Public Health, Mahidol University; 2004.
- Charurucks N. Blood lipids: Its clinical importance and laboratory method. *Chula Med J* 2006; 50: 443-58.
- Patel A, Barzi F, Jamrozik K, Lam TH, Ueshima H, Whitlock G, et al. Serum triglycerides as a risk factor for cardiovascular diseases in the Asia-Pacific region. *Circulation* 2004; 110: 2678-86.
- Jeppesen J, Hein HO, Suadicani P, Gyntelberg F. Triglyceride concentration and ischemic heart disease: an eight-year follow-up in the Copenhagen Male Study. *Circulation* 1998; 97: 1029-36.
- Hopkins PN, Heiss G, Ellison RC, Province MA, Pankow JS, Eckfeldt JH, et al. Coronary artery disease risk in familial combined hyperlipidemia and familial hypertriglyceridemia: a case-control comparison from the National Heart, Lung, and Blood Institute Family Heart Study. *Circulation* 2003; 108: 519-23.

13. Thelin A, Stiernstrom EL, Holmberg S. Blood lipid levels in a rural male population. *J Cardiovasc Risk* 2001; 8: 165-74.
14. Senti M, Aubo C, Bosch M. The relationship between smoking and triglyceride-rich lipoproteins is modulated by genetic variation in the glycoprotein IIIa gene. *Metabolism* 1998; 47: 1040-1.
15. Kuzuya M, Ando F, Iguchi A, Shimokata H. Effect of smoking habit on age-related changes in serum lipids: a cross-sectional and longitudinal analysis in a large Japanese cohort. *Atherosclerosis* 2006; 185: 183-90.
16. Hokanson JE, Austin MA. Plasma triglyceride level is a risk factor for cardiovascular disease independent of high-density lipoprotein cholesterol level: a meta-analysis of population-based prospective studies. *J Cardiovasc Risk* 1996; 3: 213-9.
17. Cullen P. Evidence that triglycerides are an independent coronary heart disease risk factor. *Am J Cardiol* 2000; 86: 943-9.
18. Abdel-Maksoud MF, Hokanson JE. The complex role of triglycerides in cardiovascular disease. *Semin Vasc Med* 2002; 2: 325-33.
19. Breuer HW. Hypertriglyceridemia: a review of clinical relevance and treatment options: focus on cerivastatin. *Curr Med Res Opin* 2001; 17: 60-73.
20. Pejic RN, Lee DT. Hypertriglyceridemia. *J Am Board Fam Med* 2006; 19: 310-6.
21. Brewer HB Jr. Hypertriglyceridemia: changes in the plasma lipoproteins associated with an increased risk of cardiovascular disease. *Am J Cardiol* 1999; 83: 3F-12F.
22. Dzien A, Dzien-Bischinger C, Hoppichler F, Lechleitner M. The metabolic syndrome as a link between smoking and cardiovascular disease. *Diabetes Obes Metab* 2004; 6: 127-32.
23. Grundy SM. Hypertriglyceridemia, insulin resistance, and the metabolic syndrome. *Am J Cardiol* 1999; 83: 25F-9F.

## ผลของการสูบบุหรี่ต่อไขมันในเลือดของประชาชนในกรุงเทพมหานคร

### ไพรัตน์ แสงดิษฐ์

**วัตถุประสงค์:** เพื่อศึกษาผลของการสูบบุหรี่ต่อระดับไขมันコレสเตอรอลและไตรกลีเซอไรด์ในเลือดในประชาชนที่จำพรรษาในกรุงเทพมหานคร

**วัสดุและวิธีการ:** ได้ตรวจระดับไขมันコレสเตอรอลและไตรกลีเซอไรด์ในการตรวจสุขภาพประชาชน 401 รายในกรุงเทพมหานครและมีการตอบแบบสอบถามเกี่ยวกับการสูบบุหรี่

**ผลการศึกษา:** 76.3 % ของประชาชนอายุมากกว่า 40 ปี อายุเฉลี่ย  $53.68 \pm 15.01$  ปี ค่าเฉลี่ยของ コレสเตอรอล ของกลุ่มที่ไม่สูบบุหรี่ กลุ่มที่เคยสูบบุหรี่แต่เลิกแล้ว กลุ่มที่สูบบุหรี่ และกลุ่มประชาชนทั้งหมดเป็น  $230.08 \pm 48.22$ ,  $223.58 \pm 42.80$ ,  $230.08 \pm 48.22$ ,  $226.65 \pm 46.58$  ตามลำดับ ค่าเฉลี่ยของไตรกลีเซอไรด์ในกลุ่มที่ไม่สูบบุหรี่ กลุ่มที่สูบบุหรี่แต่เลิกแล้ว กลุ่มที่สูบบุหรี่และประชาชนทั้งหมดเป็น  $150.05 \pm 78.23$ ,  $148.47 \pm 69.80$ ,  $180.01 \pm 111.99$ ,  $154.87 \pm 83.71$  ตามลำดับ ในการเปรียบเทียบระหว่างกลุ่มพบว่าไม่มีความแตกต่างอย่างมีนัยสำคัญทางสถิติในระดับ コレสเตอรอล ( $p = 0.22$ ) แต่มีความแตกต่างอย่างมีนัยสำคัญทางสถิติในระดับไตรกลีเซอไรด์ ( $p = 0.02$ ) เมื่อเปรียบเทียบค่าเฉลี่ยของไตรกลีเซอไรด์ระหว่างกลุ่มที่ไม่สูบบุหรี่กับกลุ่มที่สูบบุหรี่ กลุ่มที่สูบบุหรี่แต่เลิกแล้วกับกลุ่มที่สูบบุหรี่พบว่ามีความแตกต่างอย่างมีนัยสำคัญทางสถิติ ( $p = 0.026$ ,  $0.043$ ) ตามลำดับ

**สรุปผล:** จากผลการศึกษาบ่งชี้ว่าการสูบบุหรี่มีผลเพิ่มระดับไตรกลีเซอไรด์ในเลือดในประชาชน แต่ไม่มีผลต่อระดับコレสเตอรอล ดังนั้นประชาชนที่สูบบุหรี่จึงควรตรวจระดับไตรกลีเซอไรด์เพื่อจะได้มีการป้องกันการเกิดโรคหลอดเลือดหัวใจและโรคที่เกิดจากระดับไตรกลีเซอไรด์สูงอีกด้วย